

---

# Climate Change and Greenhouse Gases

---

## Authors

*Tamara S. Ledley, Eric T. Sundquist, Stephen E. Schwartz, Dorothy K. Hall, Jack D. Fellows, and Timothy L. Killeen*

For more information, contact Tamara S. Ledley, TERC, 2067 Massachusetts Ave., Cambridge, MA 02140 USA; E-mail: Tamara\_Ledley@terc.edu.

---

Copyright 1999 American Geophysical Union

---

Infrared (IR) active gases, principally water vapor (H<sub>2</sub>O), carbon dioxide (CO<sub>2</sub>), and ozone (O<sub>3</sub>), naturally present in the Earth's atmosphere, absorb thermal IR radiation emitted by the Earth's surface and atmosphere. The atmosphere is warmed by this mechanism and, in turn, emits IR radiation, with a significant portion of this energy acting to warm the surface and the lower atmosphere. As a consequence the average surface air temperature of the Earth is about 30° C higher than it would be without atmospheric absorption and reradiation of IR energy [Henderson-Sellers and Robinson, 1986; Kellogg, 1996; Peixoto and Oort, 1992].

This phenomenon is popularly known as the “greenhouse effect,” and the IR active gases responsible for the effect are likewise referred to as “greenhouse gases.” The rapid increase in concentrations of greenhouse gases since the industrial period began has given rise to concern over potential resultant climate changes.

The AGU Council approved a position statement on Climate Change and Greenhouse Gases in December 1998. The statement and a short summary of the procedures that were followed in its preparation, review, and adoption were published in the February 2, 1999, issue of *Eos* (p. 49) [AGU, 1999, also at AGU's Web site: [http://www.agu.org/sci\\_soc/policy/climate\\_change.html](http://www.agu.org/sci_soc/policy/climate_change.html)]. The present article reviews scientific understanding of this issue, as presented in peer-reviewed publications. This understanding serves as the underlying basis of the position statement.

## Greenhouse Gases and the Earth-Atmosphere Energy Balance

The principal greenhouse gas concentrations that have increased over the industrial period are carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), and chlorofluorocarbons CFC-11 (CCl<sub>3</sub>F) and CFC-12 (CCl<sub>2</sub>F<sub>2</sub>) [Hansen *et al.*, 1998; Schimel *et al.*, 1996]. The observed increase of CO<sub>2</sub> in the atmosphere from about 280 ppm in the preindustrial era to about 364 ppm in 1997 ([Figure 1](#)) [Friedli *et al.*, 1986; Hansen *et al.*, 1998; Keeling and Whorf, 1998] has come largely from fossil fuel combustion and cement production.

These sources amounted to approximately 6.5 Pg C/yr (1 Pg=10<sup>15</sup> g) in 1996 [Marland *et al.*, 1999]. Land use changes produce a non-negligible but more uncertain contribution of about  $1.6 \pm 1.0$  Pg C/yr [Fan *et al.*, 1998; Schimel *et al.*, 1996]. These anthropogenic sources of CO<sub>2</sub> exceed the estimated uptake of CO<sub>2</sub> by the atmosphere and oceans, implying a significant but as yet unidentified terrestrial sink [Enting and Pearman, 1987].

The atmospheric concentration of CH<sub>4</sub> has increased from about 700 ppb in preindustrial times to about 1721 ppb in 1994 (Figure 1) [Houghton *et al.*, 1996]. Fossil-fuel related sources of CH<sub>4</sub> amount to approximately 70-120 Tg CH<sub>4</sub>/yr (1 Tg=10<sup>12</sup> g). Increases in CH<sub>4</sub> sources resulting from rice cultivation, animal husbandry, biomass burning, and landfills contribute about 200 - 350 Tg CH<sub>4</sub>/yr [Schimel *et al.*, 1996].

The atmospheric concentration of N<sub>2</sub>O has increased from about 275 ppb in preindustrial times to about 312 ppb in 1994 (Figure 1) [Houghton *et al.*, 1996]. Estimated anthropogenic emissions of N<sub>2</sub>O for the 1980s range from 3 to 8 Tg N/yr [Houghton *et al.*, 1996]. The main anthropogenic sources are from agriculture and industrial sources including adipic acid and nitric acid production [Schimel *et al.*, 1996].

Chlorofluorocarbons CFC-12 and CFC-11 are manmade compounds that were not appreciably present in the atmosphere before 1950 (Figure 1). These compounds have been widely used as refrigerants and in spray propellants and foam blowing. Because of their role in catalyzing decomposition of stratospheric ozone, production of these compounds has been dramatically reduced in response to the Montreal Protocols and subsequent international agreements. Atmospheric concentrations of these compounds are expected to diminish substantially during the next century [Prather *et al.*, 1996].

Prediction of the future persistence of anthropogenic greenhouse gases in the atmosphere is based on mathematical models that simulate future additions and removals. The greenhouse gas concentrations predicted by these models are subject to large uncertainties in the effects of both natural processes and human activities.

For some greenhouse gases persistence can be estimated from “mean residence times,” which are obtained with simple linear models and represent the time that would be required for removal of 63% of the anthropogenic excess of the material in the atmosphere, if anthropogenic sources were abruptly diminished to zero [Lasaga and Berner, 1998]. This approach yields a rough measure of the persistence in the atmosphere of anthropogenic additions of CH<sub>4</sub> with an estimated mean residence time of 10 years [Prather, 1996, 1998]; N<sub>2</sub>O, 100 years, [Prather, 1996, 1998]; and CFC-11 and CFC-12, 50 and 102 years, respectively [Prather *et al.*, 1995].

The persistence of anthropogenic CO<sub>2</sub> in the atmosphere cannot be estimated with such a simple model because exchange with the ocean and sediments leads to a more complex behavior. Model simulations of oceanic CO<sub>2</sub> uptake provide response times associated with CO<sub>2</sub> gas exchange at the ocean surface of approximately 10 years [Liss and Merlivat, 1986; Toggweiler *et al.*, 1989] and downward mixing of surface waters on the order of decades to centuries [Maier-Reimer and Hasselmann, 1987; Sarmiento *et al.*, 1992]. But even when these oceanic CO<sub>2</sub> removal processes are allowed sufficient time in the models to reach their maximum capacity, they can remove only about 70 to 85% of the anthropogenic CO<sub>2</sub> added to the atmosphere [Archer *et al.*, 1998; Broecker and Peng, 1982; Sarmiento *et al.*, 1992].

Additional CO<sub>2</sub> might be removed by burial in soils or deep sea sediments through mechanisms that, although poorly understood, are generally believed to require times extending to thousands of years [Harden *et al.*, 1992; Schlesinger, 1990; Stallard, 1998]. Removing some of the anthropogenic CO<sub>2</sub> by this mechanism may require

reactions with carbonate sediments in the deep sea that occur on timescales of thousands of years [Archer *et al.*, 1998; Boyle, 1983; Sundquist, 1990]. On the basis of such analyses, it is now generally believed that a substantial fraction of the excess CO<sub>2</sub> in the atmosphere will remain in the atmosphere for decades to centuries, and about 15-30% will remain for thousands of years.

The additional anthropogenic greenhouse gases that have been introduced into the atmosphere increase the IR energy absorbed by the atmosphere, thereby exerting a warming influence on the lower atmosphere and the surface, and a cooling influence on the stratosphere [Peixoto and Oort, 1992; Ramanathan *et al.*, 1985].

The radiative influence resulting from a given incremental increase in greenhouse gas concentration can be quantified and compared as the change in downward IR flux at the tropopause, a quantity known as the radiative forcing. Climate model calculations indicate that to good approximation the global warming influence of the several greenhouse gases is equal for equal forcing [Wang *et al.*, 1991, 1992], lending support to the utility of the concept of climate forcing and response.

Of the several anthropogenic greenhouse gases, CO<sub>2</sub> is the most important agent of potential future climate warming because of its large current greenhouse forcing, its substantial projected future forcing [Houghton *et al.*, 1996], and its long persistence in the atmosphere (see above). Understanding climate response to a specified forcing is one of the major challenges facing the climate research community. The equilibrium response of the nonlinear climate system depends in complex ways on various feedbacks, such as changes in water vapor concentration and cloudiness that can augment or diminish climate response from that which would occur in the absence of such feedbacks.

In principle, empirical inferences of climate sensitivity would be of great value, but development of such inferences is confounded by the natural variability of the climate system [Santer *et al.*, 1996], by local or regional effects that can be different from the global effects, and by the simultaneous working of multiple transient forcings and responses. For these reasons a principal means for understanding climate system response to forcing is by use of computer models of the Earth's climate system.

## Climate Change and Carbon Dioxide

The most commonly considered indicator of climate change is the surface air temperature. Extensive efforts have been made to examine the trends in global and regional mean temperatures over time [Ghil and Vautard, 1991; Hasselmann, 1993; North and Kim, 1995; North *et al.*, 1995; Schlesinger and Ramankutty, 1994] and in the global patterns of temperature change [Hegerl *et al.*, 1997; Hegerl *et al.*, 1996; Jones and Hegerl, 1998; Santer *et al.*, 1995].

Worldwide temperature measurements, carefully screened for instrumental and measurement artifacts, such as effects of urbanization, have been used to estimate that global mean annual surface temperatures have increased between 0.3 and 0.6° C during the last 150 years [Hansen and Lebedeff, 1987; Jones *et al.*, 1997; Nicholls *et al.*, 1996]. However, it must be stressed that the increase has not been monotonic, with interannual fluctuations in the global annual mean temperature equal to an appreciable fraction of the overall rise over this time period. No single explanation can account for this variability.

Although temperature is usually the first variable considered in assessments of global climate change, it is important to consider other data that integrate the state of the climate system over space and time. These include temperature proxy data (such as tree ring records), borehole temperature measurements in soil, permafrost, and ice sheets, and measurements of the mass balance of valley glaciers and ice caps. Several recent proxy temperature reconstructions

have suggested that the warming during the twentieth century is greater than any seen in the last 400 to 600 years [Briffa *et al.*, 1998; Jones *et al.*, 1998; Mann *et al.*, 1998; Overpeck *et al.*, 1997] and perhaps the last 1200 to 1500 years (Figure 2) [Overpeck, 1998; Thompson *et al.*, 1993]. A completely independent estimate [Pollack *et al.*, 1998], based on analysis of subsurface (borehole) temperature measurements, supports the unusual character of the recent global warming in the context of the last 5 centuries.

Glaciers are present on every continent except Australia; they are thus excellent geographically dispersed regional indicators of climate change. The Earth's valley glaciers, ice caps, and ice fields and their associated outlet glaciers have generally been shrinking and receding during the last century [Haeberli, 1990; Meier, 1984; Oerlemans, 1994]. Studies in North America [Hall *et al.*, 1992; Marcus *et al.*, 1995; Rabus *et al.*, 1995; Williams and Ferrigno, in press], South America [Thompson *et al.*, 1995; Williams and Ferrigno, 1998], Europe [Williams and Ferrigno, 1993; Bayr *et al.*, 1994; Haeberli and Hoelzle, 1995], Iceland [Johannesson and Sigurðsson, 1998], Africa [Hastenrath, 1989; Williams and Ferrigno, 1991], and Asia [Thompson *et al.*, 1989, 1993, 1998] have shown substantial recession of many of the ice caps and nontidewater, nonsurge-type glaciers [Dyurgerov and Meier, 1997] since the early nineteenth century.

The record of the past few thousand years is more difficult to piece together than the more recent record because fewer data are available. There is evidence from this period that climatic conditions were sometimes warmer and sometimes cooler than at present [Dahl-Jensen *et al.*, 1998; Feng and Epstein, 1994; Prentice *et al.*, 1998].

There is also evidence of large and abrupt climate changes that exceed recent experience [Bond *et al.*, 1997; Denton and Karlen, 1973; Gasse and Van Campo, 1994; Laird *et al.*, 1996; Petitmaire and Guo, 1996; von Grafenstein *et al.*, 1998]. These climatic variations occurred during a time when variations in atmospheric CO<sub>2</sub> were minimal [Barnola *et al.*, 1995; Indermuhle *et al.*, 1999].

It is clear from these records, and from many other studies of paleoclimate evidence throughout the geologic record, that the global climate system has been influenced by many factors in addition to greenhouse gases [see, for example, Berger and Crowell, 1982]. To evaluate geologic evidence for the influence of greenhouse gases, one must focus on records from periods when changes in atmospheric CO<sub>2</sub> were much larger than those that occurred during the millennia immediately preceding the recent increase in anthropogenic CO<sub>2</sub> production.

Larger natural variations in atmospheric CO<sub>2</sub> have been inferred from the geologic record of the more distant past [for an overview, see Sundquist and Broecker, 1985]. Variations of 80-100 ppm, observed in analyses of gas bubbles trapped in glacier-ice cores, are correlated with the glacial ("ice age") and interglacial climatic oscillations of the latest Pleistocene and Holocene Epochs (Figure 3; [Barnola *et al.*, 1987; Berner *et al.*, 1980; Jouzel *et al.*, 1993; Wahlen *et al.*, 1998]). Glacial periods are associated with low CO<sub>2</sub> concentrations, and interglacial periods with high CO<sub>2</sub> concentrations. Ice core methane profiles show a similar correlation with climate [Chappellaz *et al.*, 1990; Delmotte *et al.*, 1998; Jouzel *et al.*, 1993; Stauffer *et al.*, 1988].

Still larger past variations in atmospheric CO<sub>2</sub>, including increases to concentrations several times higher than recent levels, have been estimated using geochemical models constrained by the sediment record [Berner, 1994, 1997; Berner *et al.*, 1983; Budyko and Ronov, 1979; France-Lanord and Derry, 1997; Francois and Godderis, 1998; Raymo *et al.*, 1988]. During the last several hundred million years, these larger and slower CO<sub>2</sub> changes can be correlated with general features of climate change [Berner, 1990; Crowley and North, 1991; Fischer, 1981].

Paleoclimate model simulations (using models similar in many ways to the models used in modern climate projections) support the importance of CO<sub>2</sub> in explaining global mean temperatures in the geologic past [Berger *et*

*al.*, 1998; *Bush and Philander*, 1997; *Kasting and Ackerman*, 1986; *Ottobliesner*, 1996; *Tarasov and Peltier*, 1997; *Weaver et al.*, 1998]. Model simulations have also shown the importance of changes in other climate controls, for example the configuration of Earth's orbit [*Berger and Loutre*, 1997; *Kutzbach et al.*, 1988] and the geographical distribution and elevation of continental areas [*Barron*, 1985; *Kutzbach et al.*, 1989].

Significant gaps remain in understanding the relationships among these diverse climatic influences. However, the prevailing paradigm in paleoclimate research treats the radiative effects of atmospheric CO<sub>2</sub> as an integral component in a complex system of many variables and interactive influences on global climate.

The complexity of the long-term coupling of CO<sub>2</sub> and climate is enhanced by the extent to which climate variability is hypothesized to have influenced past atmospheric CO<sub>2</sub> concentrations. Glacial/interglacial CO<sub>2</sub> variations during the Pleistocene epoch appear to have involved a combination of changes in global carbon cycling that were probably driven by some aspect of climate change [*Boyle*, 1988; *Broecker*, 1982; *Broecker and Henderson*, 1998; *Crowley*, 1995; *Heinze et al.*, 1991; *Shackleton*, 1977; *Sundquist*, 1993]. Likewise, for timescales of millions of years and longer, atmospheric CO<sub>2</sub> appears to have been affected by the influence of climate on weathering and erosion rates [*Berner*, 1990, 1994; *Berner et al.*, 1983; *Walker et al.*, 1981].

Thus current interpretation of the geologic record suggests that greenhouse gases both respond and contribute to climate change. Atmospheric CO<sub>2</sub> is viewed as one of many components of the climate system that interact in complex ways over a wide range of timescales. A change in one of these interactive components is likely to affect other aspects of the global climate system. This interactive relationship between CO<sub>2</sub> and climate implies that the geologic record is not likely to reveal analogs of simple climate forcing by anthropogenic CO<sub>2</sub> emissions [*Crowley*, 1997; *Hay et al.*, 1997; *Sundquist*, 1986]. There is no known geologic precedent for large increases of atmospheric CO<sub>2</sub> without simultaneous changes in other components of the carbon cycle and climate system.

### **Predicted Climate Change from Increased Greenhouse Gases**

Some of the predicted responses to increases in greenhouse gases include increases in mean surface air temperature, increases in global mean rates of precipitation and evaporation, rising sea level, and changes in the biosphere. Many of these predictions are based largely on computer models that simulate fundamental geophysical processes.

Most model simulations of Earth's climate indicate that an increase in the atmospheric concentration of a greenhouse gas will lead to an increase in the average surface air temperature of the Earth [*Kattenberg et al.*, 1996, table 6.3]. For example, the 18 model runs (using 7 independent models) quoted by Kattenberg et al. predict an equilibrium temperature increase of  $2.0 \pm 0.6^\circ \text{C}$  for simulations using double the current level of atmospheric CO<sub>2</sub>.

An increase in surface air temperature would cause an increase in evaporation and generally higher levels of atmospheric water vapor. The positive feedback associated with this leads to the expectation that an increase in surface air temperatures would lead to a more intense hydrological cycle, with more frequent heavy precipitation events [*Houghton et al.*, 1992; *Kattenberg et al.*, 1996]. However, because of the coarse spatial resolution of present general circulation models, simulations of the regional and seasonal distribution of precipitation are poor [*Kattenberg et al.*, 1996].

Another possible consequence of greenhouse-gas-induced climate change is elevated sea level. The main factors that contribute to sea level rise are thermal expansion of ocean water and the melting of glaciers, both of which are in response to higher air temperatures. Although it has been well established that meltwater from the world's small glaciers has contributed to sea level rise during the last century [*Dyurgerov and Meier*, 1997; *Meier*, 1984], the mass

balance of the ice sheets in Greenland and Antarctica is unknown. However, recent geodetic airborne laser altimeter measurements indicate that between 1993 and 1998 the southeastern part of the Greenland ice sheet thinned overall, with a thickening at a rate of  $0.5 \pm 0.7$  cm/yr at elevations above 2000 m (not corrected for crustal motion) and a thinning at the low elevations at rates up to 1 m/yr [Krabill *et al.*, 1999].

Worldwide measurements from tidal gauges during the last 100 years indicate that mean sea level has risen between 10 and 25 cm (18 cm mean) [Douglas, 1991, 1992; Gornitz, 1995; Warrick *et al.*, 1996]. This rate is greater than would be expected from the archaeological and geological record of sea level from the last two millennia [Warrick *et al.*, 1996]. Most modeling studies, including simulations of the combined effects of increasing greenhouse gases and aerosols, predict that the trend in rising sea level will continue in the future [Titus and Narayanan, 1995; Warrick and Oerlemans, 1990; Warrick *et al.*, 1996; Wigley and Raper, 1992, 1993].

A possible biological effect may be seen in evidence that there has been an increase in the active growing season at high latitudes in the Northern Hemisphere [Keeling *et al.*, 1996; Myneni *et al.*, 1997].

### **Predictive Capabilities and Uncertainties**

The models that have been used to study climate change are necessarily simplified representations of the climate system. Despite the inevitable limitations, climate model simulations accurately reproduce the large-scale seasonal distributions of pressure and temperature. In addition, the large-scale structure of precipitation and ocean surface heat flux also closely resembles the observed estimates [Gates *et al.*, 1998].

Confidence in models is also gained from their emerging predictive capability. An example of this capability is the development of a hierarchy of models to study the El Niño-Southern Oscillation (ENSO) phenomena [Neelin and Latif, 1998]. These models are becoming capable of predicting sea surface temperature anomalies in the tropical Pacific 6 to 12 months in advance [Latif *et al.*, 1998]. The models cannot predict specific storms related to ENSO, but they can predict the lower frequency responses of the climate system, such as anomalies in monthly and seasonal averages of the sea surface temperatures in the tropical Pacific [Neelin and Latif, 1998].

Despite these gains there are a number of features of the climate system that are still rather crudely represented in climate models. The coarse resolution of these models (typically  $3^\circ$  or roughly 300 km) restricts their ability to accurately represent terrain effects and to simulate processes that occur on smaller scales. Other shortcomings occur in the representation of aerosols, precipitation, and clouds and changes in solar irradiance. For these and other reasons there remain substantial scientific uncertainties in model predictions, including uncertainties in the predictions of local effects of climate change, occurrence of extreme weather events, effects of aerosols, changes in clouds, shifts in the intensity and distribution of precipitation, and changes in oceanic circulation [Hansen *et al.*, 1998; Houghton *et al.*, 1996; Mahlman, 1997].

A principal source of uncertainty in modeling climate change during the industrial period arises from uncertainties in the representation of the influence of anthropogenic aerosols. Aerosols scatter and absorb short wave (solar) radiation and modify the reflectivity of clouds. Both effects are thought to decrease the absorption of short wave radiation by the Earth, exerting a cooling influence on climate, despite the fact that tropospheric aerosols are short lived in the atmosphere (a few days) [Charlson *et al.*, 1992; Kaufman and Fraser, 1997; Twomey *et al.*, 1984; Haywood *et al.*, 1999].

Recent climate modeling studies which include the effects of aerosols [Hasselmann, 1997; Hegerl *et al.*, 1997; Houghton *et al.*, 1995; Kattenberg *et al.*, 1996; Mitchell *et al.*, 1995; Roeckner *et al.*, 1996] show improved comparisons between the simulated and observed global temperature trends during the industrial period. However, given the present large uncertainties in aerosol forcing, such improvement may only be fortuitous.

An additional uncertain contribution to radiative forcing of climate change during the industrial period arises from possible changes in solar irradiance. Based on reconstructions of solar irradiance and climate response in the preindustrial era, together with instrumental records and solar observations during the industrial period, Lean and Rind [1998] estimate that solar forcing may have contributed about half of the observed surface warming since 1900.

Uncertainties regarding clouds and the hydrological cycle and their representation in climate models also introduce uncertainty into present understanding of the response of the climate system to increases in atmospheric greenhouse gases. It has been indicated in model calculations that warming in the lower atmosphere as a result of greenhouse gases would increase the abundance of water vapor in the atmosphere and intensify the hydrologic cycle [*Gates et al.*, 1992; *Kattenberg et al.*, 1996].


These changes might be expected to lead to an enhancement of cloudiness. Clouds reduce the net absorbed short wave radiation in the climate system because of their high reflectivity (a cooling influence); however, they also radiate energy back down to the surface and lower atmosphere (a warming influence). The overall effect of these opposing influences is a net cooling [*Ramanathan et al.*, 1989] although this varies regionally, with cloud type, and with geography. The question of whether average cloudiness would be increased or decreased in a greenhouse-enhanced world is not yet established. Issues such as these contribute to the present uncertainty in climate sensitivity.

## Summary


In summary, the atmospheric concentrations of the principal anthropogenic greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, CFC-11, and CFC-12) have increased significantly during the industrial period. Elevated concentrations are predicted to persist in the atmosphere for times ranging to thousands of years. The increased atmospheric levels of these gases, especially CO<sub>2</sub>, increase the IR energy absorbed by the atmosphere, thereby producing a warming influence at the Earth's surface.

Global mean temperatures have increased between 0.3 and 0.6° C during the last 150 years. This change has not been monotonic, but it is unusual in the context of the last few centuries. On the timescale of the last few thousand years there have been larger climatic variation during times when variations in CO<sub>2</sub> have been relatively low. It is clear that atmospheric CO<sub>2</sub> is not the only influence on global climate. However, there have been large natural variations of CO<sub>2</sub> in the geologic past, and these changes are correlated with general features of climate change. There is no known geologic precedent for large increases of atmospheric CO<sub>2</sub> without simultaneous changes in other components of the carbon cycle and climate system.


Changes in the climate system that are confidently predicted in response to increases in greenhouse gases include increases in mean surface air temperature, increases in global mean rates of precipitation and evaporation, rising sea level, and changes in the biosphere. Substantial uncertainties remain in the magnitudes and geographical distribution of these changes and in the rates at which they may be expected to occur. The significant recent progress in the scientific understanding of climate change and the uncertainties in predictions of climate change are documented in the peer-reviewed literature. Peer-reviewed scientific research provides the scientific basis for the AGU position statement on Climate Change and Greenhouse Gases and must continue to be utilized in informed decision making on this issue.



**Fig. 1.** Concentrations of principal anthropogenic greenhouse gases in the industrial era [Hansen *et al.*, 1998; Hansen and Sato, 1999]. Black curves denote measurements of in situ atmospheric samples collected in recent years [NOAA, 1999a, b, c; Houghton *et al.*, 1995]. Points denote concentrations determined from air bubbles trapped in polar ice sheets using ice cores obtained in Antarctica (blue) or Greenland (yellow); red curves denote fits to these points [Etheridge *et al.*, 1996, 1998; Machida *et al.*, 1995]. Data for CFCs are from in situ samples since 1977 [NOAA, 1999d]. Mixing ratios of CFC-11 and CFC-12 prior to the first in situ atmospheric measurements were estimated from industrial production data and assumed atmospheric lifetimes of 50 and 100 years, respectively [AFEAS, 1993; Hansen *et al.*, 1998; Hansen and Sato, 1999].



**Fig. 2.** Reconstruction of Northern Hemisphere temperature anomaly trend from 1000 A.D. to present [*Mann et al.*, 1999; see also *Mann et al.*, 1998] from dendroclimatic, coral, and ice-core proxy records as calibrated by instrumental measurements [*Jones and Briffa*, 1992]. Thin curves give reconstruction and raw data from 1000-1998 A.D. Smoothed version (thick solid), linear trend from 1000 to 1850 A.D. (long dashed), and two standard error limits (shaded) are also shown.



**Fig. 3.** Carbon dioxide concentration (top), proxy temperature (middle), and methane concentration from analyses of ice cores from Vostok, Antarctica [Jouzel *et al.*, 1993].

## References

AGU, Climate Change and Greenhouse Gases, *Eos*, 80 (5), 49, 1999.

Alternative Fluorocarbon Environmental Acceptability Study (AFEAS), *Production, sales and atmospheric release of fluorocarbons through 1992*, Washington D.C., 1993.

Archer, D., H. Kheshgi, and E. Maier-Reimer, Dynamics of fossil fuel CO<sub>2</sub> neutralization by marine CaCO<sub>3</sub>, *Global Biogeochemical Cycles*, 12 (2), 259-276, 1998.

Barnola, J. M., M. Anklin, J. Porcheron, D. Raynaud, J. Schwander, and B. Stauffer, CO<sub>2</sub> evolution during the last millennium as recorded by Antarctic and Greenland ice, *Tellus*, 47B, 264-272, 1995.

Barnola, J. M., D. Raynaud, S. Korotkevich, and C. Lorius, Vostok ice core provides 160,000-year record of atmospheric CO<sub>2</sub>, *Nature*, 329, 408-414, 1987.

Barron, E. J., Explanations of the Tertiary global cooling trend, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 50, 45-61, 1985.

Bayr, K. J., D. K. Hall, and W. M. Kovalick, Observations on glaciers in the Eastern Austrian alps using satellite data, *International Journal of Remote Sensing*, 15 (9), 1733-1742, 1994.

Berger, A., and M. F. Loutre, Long-term variations in insolation and their effects on climate, the LLN experiments, *Surveys in Geophysics*, 18 (2-3), 147-161, 1997.

Berger, A., M. F. Loutre, and H. Gallee, Sensitivity of the LLN climate model to the astronomical and CO<sub>2</sub> forcings over the last 200 ky, *Climate Dynamics*, 14 (9), 615-629, 1998.

Berger, W. H., and J. C. Crowell, Climate in Earth history, in *Studies in Geophysics*, pp. 198, National Academy Press, Washington, D.C., 1982.

Berner, R. A., Atmospheric carbon dioxide levels over Phanerozoic time, *Science*, 249, 1382-1386, 1990.

Berner, R. A., GEOCARB II: A revised model of atmospheric CO<sub>2</sub> over Phanerozoic time, *American Journal of Science*, 294, 56-91, 1994.

Berner, R. A., The rise of plants and their effect on weathering and atmospheric CO<sub>2</sub>, *Science*, 276, 544-546, 1997.

Berner, R. A., A. C. Lasaga, and R. M. Garrels, The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years, *American Journal of Science*, 283, 641-683, 1983.

Berner, W. H., H. Oeschger, and B. Stauffer, Information on the CO<sub>2</sub> cycle from ice core studies, *Radiocarbon*, 22,

227-235, 1980.

Bond, G., W. Showers, M. Cheseby, R. Lotti, P. Almasi, P. Demenocal, P. Priore, H. Cullen, I. Hajdas, and G. Bonani, A pervasive millennial-scale cycle in North Atlantic holocene and glacial climates, *Science*, 278 (5341), 1257-1266, 1997.

Boyle, E. A., Chemical accumulation variations under the Peru current during the last 130,000 years, *J. Geophys. Res.*, 88, 7667-7680, 1983.

Boyle, E. A., Vertical oceanic nutrient fractionation and glacial/interglacial CO<sub>2</sub> cycles, *Nature*, 331 (6151), 55-56, 1988.

Briffa, K. R., P. D. Jones, F. H. Schweingruber, and T. J. Osborn, Influence of volcanic eruptions on Northern Hemisphere summer temperature over the past 600 years, *Nature*, 393 (6684), 450-455, 1998.

Broecker, W. S., Ocean chemistry during glacial time, *Geochimica et Cosmochimica Acta*, 46, 1689-1705, 1982.

Broecker, W. S., and G. M. Henderson, The sequence of events surrounding Termination II and their implications for the cause of glacial-interglacial CO<sub>2</sub> changes, *Paleoceanography*, 13 (4), 352-364, 1998.

Broecker, W. S., and T.-H. Peng, *Tracers in the Sea*, 690 pp., Eldigio Press, Palisades, New York, 1982.

Budyko, M. I., and A. B. Ronov, Chemical evolution of the atmosphere in the Phanerozoic, *Geochemistry International*, 16 (3), 1-9, 1979.

Bush, A. B. G., and S. G. H. Philander, The late Cretaceous: Simulation with a coupled atmosphere-ocean general circulation model, *Paleoceanography*, 12 (3), 495-516, 1997.

Chappellaz, J., J. M. Barnola, D. Raynaud, Y. S. Korotkevich, and C. Lorius, Ice-core record of atmospheric methane over the past 160,000 years, *Nature*, 345, 127-131, 1990.

Charlson, R. J., S. E. Schwartz, J. M. Hales, R. D. Cess, J. J. A. Coakley, J. E. Hansen, and D. J. Hofmann, Climate forcing by anthropogenic aerosols, *Science*, 255, 423-430, 1992.

Crowley, T. J., Ice age terrestrial carbon changes revisited, *Global Biogeochemical Cycles*, 9 (3), 377-389, 1995.

Crowley, T. J., The problem of paleo-analogs, *Climatic Change*, 35 (1), 119-121, 1997.

Crowley, T. J., and G. R. North, *Paleoclimatology*, 339 pp., Oxford University Press, New York, 1991.

Dahl-Jensen, D., K. Mosegaard, N. Gundestrup, G. D. Clow, S. J. Johnsen, A. W. Hansen, and N. Balling, Past temperatures directly from the Greenland Ice Sheet, *Science*, 282 (5387), 268-271, 1998.

Delmotte, M., J. Chappellaz, D. Raynaud, E. Brook, and E. Tuttle, Atmospheric methane changes during the last 400 kyr revealed by the Vostok ice core (abstract), Supplement to *Eos, Transactions, AGU*, 79 (45), F151-F152, 1998.

- Denton, G. H., and W. Karlen, Holocene climatic changes, their pattern and possible cause, *Quaternary Research*, 3, 155-205, 1973.
- Douglas, B. C., Global sea level rise, *J. Geophys. Res.*, 96 (C4), 6981-6992, 1991.
- Douglas, B. C., Global sea level acceleration, *J. Geophys. Res.*, 97 (C8), 12,699-12,706, 1992.
- Dyurgerov, M. B., and M. Meier, Mass balance of mountain and sub-polar glaciers: A new global assessment for 1961-1990, *Arctic and Alpine Research*, 29 (4), 379-391, 1997.
- Enting, I. G., and G. I. Pearman, Description of a one-dimensional carbon cycle model calibrated using techniques of constrained inversion, *Tellus*, 39B, 459-476, 1987.
- Etheridge, D. M., L. P. Steele, R. J. Francey, and R. L. Langenfelds, Atmospheric methane between 1000 A.D. and present: Evidence of anthropogenic emissions and climatic variability, *J. Geophys. Res.-Atmos.*, 103, 15,979-15,993, 1998.
- Etheridge, D., L. Steele, R. Langenfelds, R. Francey, J. Barnola, and V. Morgan, Natural and anthropogenic changes in atmospheric CO<sub>2</sub> over the last 1000 years from air in Antarctic ice and firn, *J. Geophys. Res.*, 101, 4115-4128, 1996.
- Fan, S., M. Gloor, J. Mahlman, S. Pacala, J. Sarmiento, T. Takahashi, and P. Trans, A large terrestrial carbon sink in North America implied by atmospheric and oceanic carbon dioxide data and models, *Science*, 282, 442-446, 1998.
- Feng, X. H., and S. Epstein, Climatic implications of an 8000-year hydrogen isotope time series from bristlecone pine trees, *Science*, 265 (5175), 1079-1081, 1994.
- Fischer, A. G., Climatic oscillations in the biosphere, in *Biotic Crises in Ecological and Evolutionary Time*, edited by M. H. Nitecki, pp. 103-131, Academic Press, New York, 1981.
- France-Lanord, C., and L. A. Derry, Organic carbon burial forcing of the carbon cycle from Himalayan erosion, *Nature*, 390, 65-67, 1997.
- Francois, L. M., and Y. Godderis, Isotopic constraints on the Cenozoic evolution of the carbon cycle, *Chemical Geology*, 145 (3-4), 177-212, 1998.
- Friedli, H., H. Lotscher, H. Oeschger, U. Siegenthaler, and B. Stauffer, Ice core record of <sup>13</sup>C/<sup>12</sup>C ratio of atmospheric CO<sub>2</sub> in the past two centuries, *Nature*, 324, 237-238, 1986.
- Gasse, F., and E. Vancampo, Abrupt post-glacial climate events in West Asia and North Africa monsoon domains, *Earth Planetary Science Letters*, 126 (4), 435-456, 1994.
- Gates, L., J. F. B. Mitchell, G. J. Goer, U. Cubasch, and V. P. Meleshko, Climate modeling, climate prediction and model validation, in *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment*, edited by J. T. Houghton, B. A. Callander, and S. K. Varney, pp. 97-134, Cambridge University Press, Cambridge, England, 1992.

Gates, W. L., J. S. Boyle, C. C. Covey, C. G. Dease, C. M. Doutriaux, R. S. Drach, M. Fiorino, P. J. Gleckler, J. J. Hnilo, S. M. Marlais, T. J. Phillips, G. L. Potter, B. D. Santer, K. R. Sperber, K. E. Taylor, and D. N. Williams, *An overview of the results of the atmospheric model intercomparison project (AMIP)*, UCRL-JC-129928 PCMDI Report No. 45, The Program for Climate Model Diagnosis and Intercomparison, Lawrence Livermore National Laboratory, Livermore, California, 1998.

Ghil, M., and R. Vautard, Interdecadal oscillations and the warming trend in global temperature time series, *Science*, 199, 1065-1068, 1991.

Gornitz, V., Sea Level Rise: A review of recent, past and near-future trends, *Earth Surface Processes and Landforms*, 20, 7-20, 1995.

Haeberli, W., Glacier and permafrost signals of 20th century warming, *Annals of Glaciology*, 14, 99-101, 1990.

Haeberli, W., and M. Hoelzle, Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: A pilot study with the European Alps, *Annals of Glaciology*, 21, 206-212, 1995.

Hall, D. K., J. R. S. Williams, Jr., and K. J. Bayr, Glacier recession in Iceland and Austria, *Eos*, 73 (12), 129, 135, 141, 1992.

Hansen, J., and S. Lebedeff, Global trends of measured surface air temperature, *J. Geophys. Res.*, 92, 13,345-13,372, 1987.

Hansen, J. E., M. Sato, A. Lacis, R. Ruedy, I. Gegen, and E. Matthews, Climate forcings in the industrial era, *Proceedings of the National Academy of Sciences*, 95, 12,753-12,758, 1998.

Hansen J., and M. Sato, *Well-mixed anthropogenic greenhouse gases*, <http://www.giss.nasa.gov/~cdmss/data/ghgases>, May 28, 1999.

Harden, J. W., E. T. Sundquist, R. F. Stallard, and R. K. Mark, Dynamics of soil carbon during deglaciation of the Laurentide Ice Sheet, *Science*, 258, 1921-1924, 1992.

Hasselmann, K., Optimal fingerprints for the detection of time dependent climate change, *Journal of Climate*, 6, 1957-1971, 1993.

Hasselmann, K., Climate change: Are we seeing global warming?, *Science*, 276, 914-915, 1997.

Hastenrath, S., Ice flow and mass changes of Lewis Glacier, Mount Kenya, East Africa: Observations 1974-86, Modelling, and Predictions to the Year 2000 A.D., *Journal of Glaciology*, 35 (121), 325-331, 1989.

Hay, W.W., R. M. Deconto, and C. N. Wold, Climate -- Is the past the key to the future?, *Geologische Rundschau*, 86 (2), 471-491, 1997.

Haywood, J. M., v. Ramaswamy, and B. J. Soden, Tropospheric aerosol climate forcing in clear-sky satellite observations over the oceans, *Science*, 283, 1299-1303, 1999.

Hegerl, G. C., K. Hasselmann, U. Cubasch, J. F. B. Mitchell, E. Roeckner, R. Voss, and J. Waskewitz, On multi-

- fingerprint detection and attribution of greenhouse gas and aerosol forced climatic change, *Climate Dynamics*, 13, 613-634, 1997.
- Hegerl, G. C., H. v. Storch, K. Hasselmann, B. D. Santer, U. Cubasch, and P. D. Jones, Detecting greenhouse-gas-induced climate change with an optimal fingerprint Method, *Journal of Climate*, 9, 2281-2306, 1996.
- Heinze, C., E. Maier-Reimer, and K. Winn, Glacial pCO<sub>2</sub> reduction by the world ocean: Experiments with the Hamburg carbon cycle model, *Paleoceanography*, 6 (4), 395-430, 1991.
- Henderson-Sellers, A., and P. J. Robinson, *Contemporary Climatology*, 439 pp., John Wiley Sons, Inc., New York, 1986.
- Houghton, J. T., B. A. Callander, and S. K. Varney, *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment*, Cambridge University Press, Cambridge, UK, 1992.
- Houghton, J. T., L. G. M. Filho, J. Bruce, H. Lee, B. A. Callander, E. Haites, N. Harris, and K. Maskell, Radiative forcing of climate change. In *Climate Change 1994*, pp. 231, Cambridge University Press, Cambridge, 1995.
- Houghton, J. T., L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, *Climate Change 1995: The Science of Climate Change, Intergovernmental Panel on Climate Change*, pp. 572, Cambridge University Press, Cambridge, GB, 1996.
- Indermuhle, A., T. F. Stocker, F. Joos, H. Fischer, H. J. Smith, M. Wahlen, B. Deck, D. Mastroianni, J. Tschumi, T. Blunier, R. Meyer, and B. Stauffer, Holocene carbon-cycle dynamics based on CO<sub>2</sub> trapped in ice at Taylor Dome, Antarctica, *Nature*, 398, 121-126, 1999.
- Johannesson, T., and O. Sigurðsson, Interpretation of glacier variations in Iceland 1930-1995, *Jokull*, 45, 27-33, 1998.
- Jones, P. D., K. R. Briffa, Global surface air temperature variations during the 20th century: Part 1-Spatial, temporal and seasonal details, *Holocene*, 1, 165-179, 1992.
- Jones, P. D., K. R. Briffa, T. P. Barnett, and S. F. B. Tett, High-resolution paleoclimatic records for the last millennium: Interpretation, integration and comparison with General Circulation Model control-run temperatures [Review], *Holocene*, 8 (4), 455-471, 1998.
- Jones, P. D., and G. C. Hegerl, Comparisons of two methods of removing anthropogenically related variability from the near-surface observational temperature field, *J. Geophys. Res.*, 103 (D12), 13,777-13,786, 1998.
- Jones, P. D., T. J. Osborn, and K. R. Briffa, Estimating sampling errors in large-scale temperature averages, *Journal of Climate*, 10 (10), 2548-2568, 1997.
- Jouzel, J., N. I. Barkov, J. M. Barnola, M. Bender, J. Chappellaz, C. Genthon, V. M. Kotlyakov, V. Lipenkov, C. Lorius, J. R. Petit, D. Raynaud, G. Raisbeck, C. Ritz, T. Sowers, M. Stievenard, F. Yiou, and P. Yiou, Extending the Vostok ice-core record of paleoclimate to the penultimate glacial period, *Nature*, 364, 407-412, 1993.
- Kasting, J. F., and T. P. Ackerman, Climate consequences of very high carbon dioxide levels in the Earth's early atmosphere, *Science*, 234, 1383-1385, 1986.

- Kattenberg, A., F. Giorgi, H. Grassl, G. A. Meehl, J. F. B. Mitchell, R. J. Stouffer, T. Tokioka, A. J. Weaver, and T. M. L. Wigley, Climate models - projections of future climate, in *Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 285-357, Cambridge University Press, Cambridge, UK, 1996.
- Kaufman, Y. J., and R. S. Fraser, The effect of smoke particles on clouds and climate forcing, *Science*, 277, 1636-1639, 1997.
- Keeling, C. D., J. F. S. Chin, and T. P. Whorf, Increased activity of northern vegetation inferred from atmospheric CO<sub>2</sub> measurements, *Nature*, 382, 146-149, 1996.
- Keeling, C. D., and T. P. Whorf, Atmospheric CO<sub>2</sub> records from sites in the SIO air sampling network, in *Trends: A Compendium of Data on Global Change*, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tenn., 1998.
- Kellogg, W. K., Greenhouse Effect, in *Encyclopedia of Climate and Weather*, edited by S. H. Schneider, pp. 368-371, Oxford University Press, New York, 1996.
- Krabill, W., E. Frederick, S. Manizade, C. Martin, J. Sonntag, R. Swift, R. Thomas, W. Wright, and J. Yungel, Rapid thinning of parts of the southern Greenland ice sheet, *Science*, 283, 1522-1524, 1999.
- Kutzbach, J. E., P. J. Guetter, W. F. Ruddiman, and W. L. Prell, Sensitivity of climate to late Cenozoic uplift in southern Asia and the American west: numerical experiments, *J. Geophys. Res.*, 94 (D15), 18,393-18,407, 1989.
- Kutzbach, J. E., T. Webb, III, H. E. Wright, Jr., and C. Members, Climatic changes of the last 18,000 years: Observations and model simulations, *Science*, 241, 1043-1052, 1988.
- Laird, K. R., S. C. Fritz, K. A. Maasch, and B. F. Cumming, Greater drought intensity and frequency before AD 1200 in the Northern Great Plains, USA, *Nature*, 384 (6609), 552-554, 1996.
- Lasaga, A. C., and R. A. Berner, Fundamental aspects of quantitative models for geochemical cycles, *Chemical Geology*, 145 (3-4), 161-175, 1998.
- Latif, M., D. Anderson, T. Barnett, M. Cane, R. Kleeman, A. Leetmaa, J. O'Brien, A. Rosati, and E. Schneider, A review of the predictability and prediction of ENSO, *J. Geophys. Res.-Oceans*, 103 (C7), 14,375-14,393, 1998.
- Lean, J., and D. Rind, Climate forcing by changing solar radiation, *Journal of Climate*, 11, 3069-3094, 1998.
- Liss, P., and L. Merlivat, Air-sea gas exchange rates, introduction and synthesis, in *The role of air-sea exchange in geochemical cycling*, edited by P. Buat-Menard, pp. 113-128, D. Reidel, Hingham, Mass., 1986.
- Machida, T., T. Nakazawa, Y. Fujii, S. Aoke, and O. Watanabe, Increase in atmospheric nitrous oxide concentrations during the last 250 years, *Geophysical Research Letters*, 22, 2921-2924, 1995.
- Mahlman, J. D., Uncertainties in projections of human-caused climate warming, *Science*, 278, 1416-1417, 1997.

- Maier-Reimer, E., and K. Hasselmann, Transport and storage of CO<sub>2</sub> in the ocean -- an inorganic ocean-circulation carbon cycle model, *Climate Dynamics*, 2, 63-90, 1987.
- Mann, M. E., R. S. Bradley, and M. K. Hughes, Global-scale temperature patterns and climate forcing over the past six centuries, *Nature*, 392 (6678), 779-787, 1998.
- Mann, M. E., R. S. Bradley, and M. K. Hughes, Northern hemisphere temperatures during the past millennium: Inferences, uncertainties, and limitations, *Geophysical Research Letters*, 26, 759-762, 1999.
- Marcus, M., F. B. Chambers, M. M. Miller, and M. Lang, Recent trends in Lemon Creek Glacier, Alaska, *Physical Geography*, 16 (2), 15-161, 1995.
- Marland, G., R. J. Andres, T. A. Boden, C. Johnson, and A. Brenkert, Global, regional, and national CO<sub>2</sub> emission estimates from fossil fuel burning, cement production, and gas flaring: 1751-1996, CDIAC, 1999.
- Meier, M. F., Contribution of small glaciers to global sea level, *Science*, 226, 1418-1421, 1984.
- Mitchell, J. F. B., T. C. Johns, J. M. Gregory, and S. F. B. Tett, Climate response to increasing greenhouse gases and sulphate aerosols, *Nature*, 376, 501-504, 1995.
- Myneni, R. B., C. D. Keeling, C. J. Tucker, G. Asrar, and R. R. Nemani, Increased plant growth in the northern high latitudes from 1981 to 1991, *Nature*, 386, 698-702, 1997.
- Neelin, J.D., and M. Latif, El Niño Dynamics, *Physics Today*, 51 (12), 32-36, 1998.
- Nicholls, N., G. V. Gruza, J. Jouzel, T. R. Karl, L. A. Ogallo, and D. E. Parker, Observed climate variability and change, in *Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 133-192, Cambridge University Press, Cambridge, UK, 1996.
- NOAA Climate Monitoring and Diagnostics Laboratory, Carbon Cycle Group (1999a) Continuous in-situ CO<sub>2</sub> data files (<ftp://ftp.cmdl.noaa.gov/ccg/co2/in-situ>).
- NOAA Climate Monitoring and Diagnostics Laboratory, Carbon Cycle Group (1999b) Atmospheric methane mixing ratios (<ftp://ftp.cmdl.noaa.gov/ccg/ch4/flask>).
- NOAA Climate Monitoring and Diagnostics Laboratory, Halocarbons Other Atmospheric Trace Species Group (1999c) Group data (<ftp://ftp.cmdl.noaa.gov/noah/n2o>).
- NOAA Climate Monitoring and Diagnostics Laboratory, Halocarbons Other Atmospheric Trace Species Group (1999d) Group data (<ftp://ftp.cmdl.noaa.gov/noah/cfcs>).
- North, G. R., and K.-Y. Kim, Detection of forced climate signals. Part II: Simulation results, *Journal of Climate*, 6, 409-417, 1995.
- North, G. R., K.-Y. Kim, S. P. Shen, and J. W. Hardin, Detection of forced climate signals. Part I: filter theory, *Journal of Climate*, 6, 401-408, 1995.

- Oerlemans, J., Quantifying global warming from the retreat of glaciers, *Science*, 264, 243-246, 1994.
- Ottobliesner, B. L., Initiation of a continental ice sheet in a global climate model (GENESIS), *J. Geophys. Res.-Atmos.*, 101 (D12), 16,909-16,920, 1996.
- Overpeck, J., K. Hughen, D. Hardy, R. Bradley, R. Case, M. Douglas, B. Finney, K. Gajewski, G. Facoby, A. Jennings, S. Lamoureaux, A. Lasca, G. MacDonald, J. Moore, M. Retelle, S. Smith, A. Wolfe, and G. Zielinski, Arctic environmental change of the last four centuries, *Science*, 278, 1251-1256, 1997.
- Overpeck, J. T., How unprecedented is recent Arctic warming: A look back to the Medieval Warm Period (abstract), Supplement to *Eos, Transactions*, 79 (45), F833-F834, 1998.
- Peixoto, J. P., and A. H. Oort, *Physics of Climate*, 520 pp., American Institute of Physics, New York, 1992.
- Petitmaire, N., and Z. T. Guo, In-phase Holocene climate variations in the present-day desert areas of China and northern Africa, *Comptes Rendus de l'Academie des Sciences Série II Fascicule A-Sciences de la Terre et des Planètes*, 322 (10), 847-851, 1996.
- Pollack, H. N., S. Huang, and P.-Y. Shen, Climate change record in subsurface temperatures: A global perspective, *Science*, 282, 279-281, 1998.
- Prather, M., R. Derwent, D. Ehhalt, P. Fraser, E. Sanhuea, and X. Zhou, Other trace gases and atmospheric chemistry, in Radiative Forcing of Climate Change, in *Climate Change 1994*, edited by J. T. Houghton, L. G. M. Filho, J. Bruce, H. Lee, B. A. Callander, E. Haites, N. Harris, and K. Maskell, pp. 72-126, Cambridge University Press, Cambridge, 1995.
- Prather, M., P. Midgley, F. S. Rowland, and R. Stolarski, The ozone layer: The road not taken, *Nature*, 381, 551-554, 1996.
- Prather, M. J., Time scales in atmospheric chemistry: Theory, GWP's for CH and CO, and runaway growth, *Geophysical Research Letters*, 23 (19), 2597-2600, 1996.
- Prather, M. J., Time scales in atmospheric chemistry: Coupled perturbations to N<sub>2</sub>O, NO<sub>y</sub>, and O<sub>3</sub>, *Science*, 279, 1339-1341, 1998.
- Prentice, I. C., S. P. Harrison, D. Jolly, and J. Guiot, The climate and biomes of Europe at 6000 yr BP: Comparison of model simulations and pollen-based reconstructions, *Quaternary Science Reviews*, 17 (6-7), 659-668, 1998.
- Rabus, B., K. Echelmeyer, D. Trabant, and C. Benson, Recent change of McCall Glacier, Alaska, *Annals of Glaciology*, 21, 231-239, 1995.
- Ramanathan, V., R. D. Cess, E. F. Harrison, P. Minnis, B. R. Barkstrom, E. Ahmad, and D. Hartmann, Cloud-radiative forcing and climate: Results from the Earth radiation budget experiment, *Science*, 243, 57-63, 1989.
- Ramanathan, V., R. J. Cicerone, H. B. Singh, and J. T. Kiehl, Trace gas trends and their potential role in climate change, *J. Geophys. Res.*, 90, 5547-5566, 1985.

- Raymo, M. E., W. F. Ruddiman, and P. N. Froelich, Influence of late Cenozoic mountain building on ocean geochemical cycles, *Geology*, 16, 649-653, 1988.
- Roeckner, E., J. M. Oberhuber, A. Bacher, M. Christoph, and I. Kirchner, ENSO variability and atmospheric response in a global coupled atmosphere-ocean GCM, *Climate Dynamics*, 12, 737-754, 1996.
- Santer, B. D., K. E. Taylor, T. M. L. Wigley, J. E. Penner, P. D. Jones, and U. Cubasch, Towards the detection and attribution of an anthropogenic effect on climate, *Climate Dynamics*, 12, 79-100, 1995.
- Santer, B. D., T. M. L. Wigley, T. P. Barnett, and E. Anyamba, Detection of climate change and attribution of causes, in *Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 407-443, Cambridge University Press, Cambridge, UK, 1996.
- Sarmiento, J. L., J. C. Orr, and U. Siegenthaler, A perturbation simulation of CO<sub>2</sub> uptake in an ocean general circulation model, *J. Geophys. Res.*, 97 (C3), 3621-3645, 1992.
- Schimel, D., D. Alves, I. Enting, M. Heimann, F. Joos, D. Raynaud, T. Wigley, M. Prather, R. Derwent, D. Ehhalt, P. Fraser, E. Sauhueza, X. Zhou, P. Jonas, R. Charlson, H. Rodhe, S. Sadasivan, K. P. Shine, Y. Fouquart, V. Ramaswamy, S. Solomon, J. Srinivasan, D. Albritton, R. Derwent, I. Isaksen, M. Lal, and D. Wuebbles, Radiative forcing of climate change, in *Climate Change 1995: The Science of Climate Change*, edited by J. T. Houghton, L. G. M. Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, pp. 65-131, Cambridge University Press, Cambridge, UK, 1996.
- Schlesinger, M. E., and N. Ramankutty, An oscillation in the global climate system of period 65-70 years, *Nature*, 360, 330-333, 1994.
- Schlesinger, W. H., Evidence from chronosequence studies for a low carbon-storage potential of soils, *Nature*, 348, 232-234, 1990.
- Shackleton, N. J., Carbon-13 in *Uvigerina*: Tropical rainforest history and the equatorial Pacific carbonate dissolution cycles, in *The Fate of Fossil Fuel CO<sub>2</sub> in the Oceans*, edited by N. R. Andersen, and A. Malahoff, pp. 401-427, Plenum Press, New York, 1977.
- Stallard, R. F., Terrestrial sedimentation and the carbon cycle: Coupling weathering and erosion to carbon burial, *Global Biogeochemical Cycles*, 12 (2), 231-257, 1998.
- Stauffer, B., E. Lochbronner, H. Oeschger, and J. Schwander, Methane concentration in the glacial atmosphere was only half that of the preindustrial Holocene, *Nature*, 332, 812-814, 1988.
- Sundquist, E. T., Geologic analogs: Their value and limitations in carbon dioxide research, in *The Changing Carbon Cycle, A Global Analysis*, edited by J. R. Trabalka, and D. E. Reichle, pp. 371-402, Springer-Verlag, New York, 1986.
- Sundquist, E. T., Long-term aspects of future atmospheric CO<sub>2</sub> and sea-level changes, in *Sea-Level Change*, edited by R. Revelle, pp. 193-207, National Academy Press, Washington, DC, 1990.
- Sundquist, E. T., The global carbon dioxide budget, *Science*, 259, 934-941, 1993.

- Sundquist, E. T., and W. S. Broecker, The carbon cycle and atmospheric CO<sub>2</sub>: Natural variations Archean to present, in *Geophysical Monograph*, pp. 627, American Geophysical Union, Washington, DC, 1985.
- Tarasov, L., and W. R. Peltier, Terminating the 100 kyr ice age cycle, *J. Geophys. Res.- Atmos.*, 102 (D18), 21,665-21,693, 1997.
- Thompson, L. G., M. E. Davis, E. Mosely-Thompson, T. A. Sowers, K. A. Henderson, V. S. Zagorodnov, P.-N. Lin, V. N. Mikhalenko, R. K. Campen, J. F. Bolzan, J. Cole-Dai, and B. Francou, A 25,000-year tropical climate history from Bolivian ice cores, *Science*, 282, 1858-1864, 1998.
- Thompson, L. G., E. Mosely-Thompson, M. Davis, P. N. Lin, T. Yao, M. Dyurgerov, and J. Dai, "Recent Warming": Ice core evidence from tropical ice cores with emphasis on central Asia, *Global and Planetary Change*, 7, 145-156, 1993.
- Thompson, L. G., E. Mosely-Thompson, M. E. Davis, J. F. Bolzan, J. Dai, T. Yao, N. Gundestrup, X. Wu, L. Klein, and Z. Xie, Holocene-late pleistocene climatic ice core records from Qinghai-Tibetan Plateau, *Science*, 246, 474-477, 1989.
- Thompson, L. G., E. Mosely-Thompson, M. E. Davis, P.-N. Lin, K. A. Henderson, J. Cole-Dai, J. F. Bolzon, and K.-B. Liu, Late glacial stage and holocene tropical ice core records from Huascarán, Peru, *Science*, 269, 46-48, 1995.
- Titus, J. G., and V. Narayanan, *The probability of sea level rise*, U.S. Environmental Protection Agency, Office of Policy, Planning, and Evaluation, Climate Change Division, Washington, D.C., 1995.
- Toggweiler, J. R., K. Dixon, and K. Bryan, Simulations of radiocarbon in a coarse-resolution, world ocean model, II, Distributions of bomb-produced <sup>14</sup>C, *J. Geophys. Res.*, 94, 8243-8264, 1989.
- Twomey, S. A., M. Piegras, and T. L. Wolfe, An assessment of the impact of pollution on global cloud albedo, *Tellus*, 36B, 356-366, 1984.
- von Grafenstein, U., H. Erlenkeuser, J. Muller, J. Jouzel, and S. Johnsen, The cold event 8200 years ago documented in oxygen isotope records of precipitation in Europe and Greenland, *Climate Dynamics*, 14 (2), 73-81, 1998.
- Wahlen, M., D. Mastroianni, B. Deck, H. Fischer, H. J. Smith, and J. Jouzel, Carbon dioxide in the Vostok ice core (abstract), Supplement to *Eos, Transactions, AGU*, 79 (45), F150, 1998.
- Walker, J. C. G., P. B. Hays, and J. F. Kasting, A negative feedback mechanism for the long-term stabilization of Earth's surface temperature, *J. Geophys. Res.*, 86 (C10), 9776-9782, 1981.
- Wang, W.-C., M. P. Dudek, and X.-Z. Liang, Inadequacy of effective CO<sub>2</sub> as a proxy in assessing the regional climate change due to other radiatively active gases, *Geophysical Research Letters*, 19, 1375-1378, 1992.
- Wang, W.-C., M. P. Dudek, X.-Z. Liang, and J. T. Kiehl, Inadequacy of effective CO<sub>2</sub> as a proxy in simulating the greenhouse effect of other radiatively active gases, *Nature*, 350, 573-577, 1991.

Warrick, R. A., and J. Oerlemans, Sea level rise, in *Climate Change, The IPCC Assessment*, edited by J. T. Houghton, G. J. Jenkins, and J. J. Ephraums, Cambridge University Press, Cambridge, UK, 1990.

Warrick, R. A., C. L. Provost, M. F. Meier, J. Oerlemans, and P. L. Woodworth, Changes in sea level, *Climate Change 1995: The Science of Climate Change*, 359-405, 1996.

Weaver, A. J., M. Eby, F. F. Augustus, and E. C. Wiebe, Simulated influence of carbon dioxide, orbital forcing and ice sheets on the climate of the Last Glacial Maximum, *Nature*, 394, 847-853, 1998.

Wigley, T. M. L., and S. C. B. Raper, Implications for climate and sea level of revised IPCC emissions scenarios, *Nature*, 357, 293-300, 1992.

Wigley, T. M. L., and S. C. B. Raper, Future changes in global mean temperature and sea level, in *Climate and Sea Level: Observations, Projections and Implications*, edited by R. A. Warrick, E. M. Barrow, and T. M. L. Wigley, pp. 111-133, Cambridge University Press, Cambridge, UK, 1993.

Williams, R. S., Jr. and J. G. Ferrigno (editors), *Satellite image atlas of glaciers of the world: U.S. Geological Survey Professional Paper 1386-G (Glaciers of the Middle East and Africa)*, 70pp., 1991.

Williams, R. S., Jr. and J. G. Ferrigno (editors), *Satellite image atlas of glaciers of the world: U.S. Geological Survey Professional Paper 1386-E (Glaciers of Europe)*, 164pp., 1993.

Williams, R. S., Jr. and J. G. Ferrigno (editors), *Satellite image atlas of glaciers of the world: U.S. Geological Survey Professional Paper 1386-I (Glaciers of South America)*, URL address: <http://pubs.usgs.gov/prof/p1386i/>, 206pp., 1998.

Williams, R. S., Jr. and J. G. Ferrigno (editors), *Satellite image atlas of glaciers of the world: U.S. Geological Survey Professional Paper 1386-J (Glaciers of North America)*, 164pp., (in press).

